

1 **A dataset of 40'000 trees with section-wise measured stem diameter**  
2 **and branch volume from across Switzerland**

3 Markus Didion<sup>1</sup>, Anne Herold<sup>1</sup>, Esther Thürig<sup>1</sup>, Serra Topuz<sup>1</sup>, Zeljka Vulovic<sup>1</sup>, Meinrad Abegg<sup>1</sup>,  
4 Jens Nitzsche<sup>1</sup>, Jonas Stillhard<sup>1</sup>, Jonas Glatthorn<sup>1</sup>

5

6 Affiliations

7 1. Forest Resources and Management, Swiss Federal Institute for Forest, Snow and Landscape  
8 Research WSL

9 corresponding author: Markus Didion (markus.didion@wsl.ch)

10

11 **Abstract**

12 Estimating growing stock is one of the main objectives of forest inventories. It refers to the  
13 stem volume of individual trees which is typically derived by models as it cannot be easily  
14 measured directly. These models are thus based on measurable tree dimensions and their  
15 parameterization depends on the available empirical data. Historically, such data were  
16 collected by measurements of tree stem sizes, which is very time- and cost-intensive.

17 Here, we present an exceptionally large dataset with section-wise stem measurements on  
18 40'349 felled individual trees collected on plots of the Experimental Forest Management  
19 project. It is a revised and expanded version of previously unpublished data and contains the  
20 empirically derived coarse (diameter  $\geq 7$  cm) and fine branch volume of 27'297 and 18'980,  
21 respectively, individual trees. The data were collected between 1888 and 1974 across  
22 Switzerland covering a large topographic gradient and a diverse species range and can thus  
23 support estimations and verification of volume functions also outside Switzerland including  
24 the derivation of whole tree volume in a consistent manner.

25 **Background & Summary**

26 Forests are an important global resource and information on forests has been collected in  
27 inventory systems for decades or centuries following country-specific approaches<sup>1</sup>. The  
28 growing stock comprising the volume of stems of standing trees over a specific forest area<sup>2</sup>,  
29 is, recognised as one of the most important variables in forest inventories, particularly in  
30 Europe<sup>3,4</sup>. Growing stock serves as an indicator of forest functions<sup>4</sup>, as the basis for the  
31 development of forest management practices<sup>5</sup>, policy making<sup>6</sup>, and international reporting<sup>7</sup>.  
32 Data on individual stems is used to study, for example, tree taper<sup>8</sup>, volume<sup>9</sup>, and growth<sup>10</sup>.

33 Growing stock and in particular the volume of individual trees cannot be measured directly.  
34 Tree volume is thus usually estimated using models developed on the basis of tree attributes  
35 that can be measured in the field. These typically include DBH, and in some countries a  
36 diameter at a second height, and total tree height<sup>4,11</sup>. Volume models are, however, generally  
37 developed based primarily on local data that are not representative on a national scale and of

38 the occurring tree species<sup>12</sup>. A reason for this is that representative large-scale sampling is  
39 typically too time-consuming and costly<sup>11</sup>. While methods for estimating the volume of stems  
40 have been developed accounting for these limitations, this is much less the case for branch  
41 volume and crown mass. Since growing stock in European forest inventories excludes the  
42 stump and branches<sup>2</sup>, it underestimates above-ground tree volume. To also account for  
43 branches additional measurements are needed. Branch volume is typically estimated using  
44 separate functions or expansion factors<sup>4</sup>. These are generally derived from independent data  
45 based on different and typically limited population samples<sup>13</sup>. Making existing datasets  
46 available to the scientific community has the potential to significantly contribute to further  
47 develop existing methods.

48 The Swiss National forest inventory (NFI) is the main source of nationally representative  
49 information on the state and change of forest volume, biomass, and carbon stocks in  
50 Switzerland. Data from the NFI are the basis for several research, monitoring, and reporting  
51 programs such as national and international forest reports and greenhouse gas reporting. In  
52 addition to the classic assessment of growing stock, bias-free and accurate estimates of whole-  
53 tree biomass and carbon stocks are therefore required. The methods applied in the Swiss NFI  
54 are continuously improved and regularly documented e.g., Brassel and Lischke<sup>14</sup>, Fischer and  
55 Traub<sup>15</sup>. The volume of above-ground coarse (i.e.  $\geq 7$  cm in diameter, including tree stump)  
56 and fine woody parts of stem and branches is estimated using functions fitted to data collected  
57 on sites of the Experimental Forest Management project's (EFM) long-term growth and yield  
58 plot network<sup>16</sup>. The EFM project collects growth and yield data in Switzerland since the late  
59 1880's on more than 1000 plots<sup>17,18</sup>. In addition to monitoring data of standing living trees,  
60 detailed measurements of felled trees were conducted in the past. The EFM is an ongoing  
61 project and long-term consistency is assured.

62 This data paper presents an exceptionally large dataset with measurements of individual trees  
63 combining stem size (diameter and length) with the volume of coarse (converted from  
64 measured size) and fine (converted from measured weight) branches. This dataset differs from  
65 the one used to derive volume functions for stem- and branchwood in the Swiss NFI<sup>16,19</sup> in that  
66 the previously separate stem- and branchwood datasets are linked at the level of individual  
67 trees. The dataset also includes additional tree measurements and metadata. By linking stem-  
68 and branchwood measurements for individual trees, a consistent total above-ground tree  
69 volume can be derived. This information can be used to evaluate the accuracy of typical  
70 approaches to obtain total tree volume, such as adding up estimates based on two separate  
71 models or applying expansion factors which are based on different tree populations. The data  
72 can be used to further develop existing volume estimates in the Swiss NFI, resulting in higher  
73 accuracy of derived variables such as biomass and C stocks. Open access to the dataset can  
74 also support the estimation and verification of volume functions also outside Switzerland, as  
75 growing stock and total tree biomass are among the most important variables in forest  
76 inventories<sup>4</sup>.

77

## 78 **Methods**

79 Starting in the 1880s, the EFM project is one of the longest running scientific projects in  
80 Switzerland with the primary objective to provide long-term empirical data to examine forest  
81 development under the influence of management and changing environmental conditions<sup>17</sup>.  
82 Besides repeated measurements on living trees, comprehensive measurements on felled trees

83 have also been conducted throughout the years, which are the object of this data paper.  
84 Individual tree data for a range of tree species (Table 1) were obtained following the field  
85 procedure described in Flury<sup>20,21</sup>. First, all trees were numbered and marked at the height of  
86 1.30 m where the DBH was measured crosswise. Trees to be felled were selected based on  
87 Ulrich's method<sup>22,23</sup>. On a subset of the felled trees, detailed length and diameter  
88 measurements on coarse stem and branch parts as well as weights of fine parts were obtained  
89 (see section 'Data records' and Table 2).

90 The coarse woody part of the stem starting from the base of a tree up to the diameter  
91 threshold of 7 cm (i.e. including stump and bole following Gschwantner, et al.<sup>24</sup>) was divided  
92 into sections of 2 m length and the diameter at half the length of each section was measured  
93 crosswise (Figure 1). The section-wise diameter measurements therefore started at 1m from  
94 the tree base and were continued along the stem up to the thinner end where the diameter  
95 was 7 cm. As the final section was considered where the coarse stemwood diameter reached  
96 the lower bound of 7 cm. If the length of the final section was < 2 m, its full length and diameter  
97 at half its length were measured. On a subset of the trees an additional diameter  
98 measurement was made at 0.65 m. Section-wise measurements were also made on coarse  
99 branches but based on a section length of 1m. The coarse branch volume was calculated based  
100 on the section length and diameter at half the section length.

101 The parts of stem and branches below the diameter threshold of 7 cm (henceforth stem top  
102 and fine branches, respectively) were collected and fitted into standardised bundles of 1m  
103 length and 1m circumference. The fresh weight of bundles was measured directly in the field.  
104 for conversion to standardised volume<sup>25</sup>. Conversion factors (Table 3) were used to calculate  
105 the volume of standardized bundles from their fresh weight<sup>25</sup>. These were derived from data  
106 collected in the years 1888 to 1892. The data comprised both, measured fresh weight and  
107 xylometric volume for a total of 2192 standardized bundles with fine woody material collected  
108 on a representative subset of the EFM sites for the tree species *Picea*, *Abies*, *Pinus*, *Fagus*, and  
109 *Fraxinus*. The conversion factors derived by Flury<sup>25</sup> were reviewed in 1940 and expanded with  
110 more precise data for additional species. Revised factors (Table 3) were based on data from  
111 Gayer and Fabricius<sup>26</sup>. Since the factors after Gayer and Fabricius<sup>26</sup> were developed for  
112 stemwood, values were slightly modified based on expert knowledge for the application to  
113 fine woody material. The revised factors were applied for weight to volume conversion  
114 starting 1940.

115 All field measurements were recorded on paper copies of field record forms. The documents  
116 are available in the research collection of the WSL archive under "Wissenschaftliche Sammlung  
117 Ertragskunde" and partially also uncatalogued in the EFM archive. In 1974 data on measured  
118 stem dimensions from the field recording forms were converted to punchcards and over time  
119 they were also converted to a digital format. Branchwood data were processed in a separate  
120 project in 1984. This resulted in two independent datasets, one for stem dimensions  
121 (N=38'864 individual tree data) and one for branch volume (N=14'712). These datasets were  
122 the basis for the existing volume models in the Swiss NFI<sup>16,19</sup>. Documentation of this work is  
123 available on handwritten notes, and for the branchwood data in 1984 also in a detailed project  
124 proposal. Due to missing metadata, it was not straightforward to recognize whether a  
125 correlation between the two datasets existed. The here presented dataset (henceforth current  
126 dataset) is the result of research on the provenance of the initial separate stem and branch  
127 datasets that allowed to link the measured data for individual trees. Furthermore, to extend  
128 the DBH and elevation range as well as to increase the sample size of trees from uneven-aged

129 forests, the current dataset was expanded by digitizing additional tree data from the original  
130 paper copies.

## 131 **Data Records**

132 The current dataset contains measurements of 40'349 individual trees collected on 768 EFM  
133 sample plots (Figure 2). All available variables including their units and summary statistics are  
134 presented in Table 2. Figure 3 shows the DBH distribution, Figure 4 the diameter distributions  
135 of different stem sections, and Figure 5 the calculated volume of coarse and fine branches in  
136 relation to tree DBH. The dataset covers information from 768 plots (Figure 2), excluding  
137 subplots, collected at variable intervals in the period 1888 to 1974. It includes latitude,  
138 longitude and elevation of the plot centre. Site identifiers for each record can be used to derive  
139 further site metadata<sup>18</sup>. The tree data include information on

- 140 • tree species (N=28),
- 141 • tree age (based on year ring count; mean 73 years),
- 142 • DBH (mean 255 mm; Figure 3)
- 143 • total length of the stem (i.e. tree height; mean=219 dm),
- 144 • length of the coarse stemwood (from the base of a tree to the diameter threshold of  
145 7 cm (mean 180 dm),
- 146 • length of the final section of the coarse stemwood (mean 7 dm),
- 147 • length of the tree top (i.e. starting where the stem diameter is 7 cm to the end of the  
148 stem; mean=39 dm),
- 149 • mean of crosswise measured diameters over bark along the stem at 0.65 m and every  
150 2 m starting at 1m from the tree base up to the length of the coarse stemwood (Figure  
151 4)
- 152 • diameter of the final section if less than 2 m at half its length (mean 48 mm),
- 153 • diameter of the tree top (part of the stem where  $D < 7$  cm measured at half its length  
154 (mean 39 mm),
- 155 • volume of coarse branches (derived from the measured diameters at the middle of  
156 one-meter sections(mean 16 dm<sup>3</sup>; Figure 5),
- 157 • volume of all fine woody parts (ie. fine branches and tree tops derived from the  
158 measured weight of standardized bundles (mean 210 dm<sup>3</sup>; Figure 5), and
- 159 • volume of fine branches (where measured separately; derived from the measured  
160 weight of standardized bundles mean 141 dm<sup>3</sup>).

161 The quality controlled (see section 'Technical Validation') data are stored in table format as  
162 comma-separated file (.csv). The file is available from the environmental data portal EnviDat  
163 of the Swiss Federal Institute for Forest, Snow and Landscape Research WSL  
164 (<https://www.doi.org/10.16904/envi.dat.486>). Missing values or not measured variables are  
165 denoted by NA. Values of '0' indicate true values, e.g. in the case of coarse branchwood on  
166 spruce (*P. abies*) trees that generally only possess fine branches<sup>27</sup>.

## 167 **Technical Validation**

168 In a first step, the two initially available and separate digital datasets were assessed for  
169 consistency with field records and plausibility. The examination of the field recording forms  
170 also allowed the tree measurement procedure to be confirmed. Detailed information on the  
171 field procedure with cross-reference to field recording forms are available in section 3.2 in

172 Didion, et al. <sup>28</sup>. The plausibility of tree attribute values was evaluated using consistency checks  
173 to identify, for example, duplicate tree records, cases where the diameter of stem sections  
174 increased from the base to the top of the stem, or where tree height was less than the length  
175 of the merchantable part of the stem. Outlier detection was used to examine values of  
176 individual variables and in combination, for example the height to DBH ratio (Figure 6), and  
177 diameters along the stem. The quality control and merge of the stem and branch datasets was  
178 achieved in several successive steps making use of the common variables, i.e. site information,  
179 inventory year, tree species, DBH, diameter at 7 m, and total height. The correct merge by  
180 individual trees was verified by comparing with field recording forms.

181

## 182 **Usage Notes**

183 Although this dataset with consistent and detailed measurements of nearly 40'000 individual  
184 trees is very comprehensive, it should be noted that:

- 185 • particularly the Swiss regions of the Southern and Western Alps are not well  
186 represented with only few sites in the Valais and none in Ticino (Figure 1);
- 187 • mountain forest at higher elevations (> 1500 m) are poorly covered in comparison to  
188 the forest distribution based on the Swiss NFI;
- 189 • the majority of the data comes from homogenous, even-aged forests.

190 The dataset provides an empirically derived stem and branch volume. It can be used to  
191 calibrate allometric functions with variables that are easy to measure in the field such as DBH,  
192 tree height as well as a second diameter. The comprehensive dataset can also be used to  
193 examine alternative stem volume estimations based on, for example, a cylindrical first section  
194 and adding truncated cones using the mid-diameters of further sections to represent top and  
195 bottom, or taper functions<sup>11,29</sup>.

## 196 **Code Availability**

197 All data processing including quality controls and figure generation was done using the  
198 language and environment for statistical computing R version 4.2.1<sup>30</sup> and the packages  
199 data.table<sup>31</sup>, ggplot2<sup>32</sup>, and dplyr<sup>33</sup>.

## 200 **Acknowledgements**

201 We are grateful to J. Zell (formerly WSL) for discussion, A. Zingg, retired EFM lead, for valuable  
202 advice and to C. Hoffmann, retired WSL statistician, for answering our questions on his project  
203 in the mid-1980s as well as to E. Kaufmann, retired WSL staff, for his work with the initial  
204 datasets. We also thank A. Zurlinden, archive coordinator at WSL, for his help to locate  
205 documents in the WSL archive. The preliminary data explorations of S. Liechti provided a  
206 valuable basis for the further review and evaluation of the stem and branch datasets. Part of  
207 this work was supported by the Swiss Federal Office for the Environment.

## 208 **Author contributions**

209 MD: primary author, derivation of data origin and history, data matching, plausibility checks  
210 and processing.

211 AH: derivation of data origin and history, data matching and plausibility checks.

212 ZV, JN, JS, JG: data archive operation and maintenance, EFM expertise - derivation of data  
213 origin and history, review.

214 ET, MA: project support, review

215 ST: data digitalization

216 All authors contributed to the manuscript text.

## 217 **Competing interests**

218 The authors declare no competing interests.

## 219 **References**

220 1 Tomppo, E., Gschwantner, T., Lawrence, M. & McRoberts, R. E. 612 (Springer,  
221 Heidelberg, 2010).

222 2 Lanz, A. *et al.* A Sample of COST Action E43 Reference Definitions in *National forest*  
223 *inventories : pathways for common reporting* (eds E. Tomppo, Th Gschwantner, M.  
224 Lawrence, & R.E. McRoberts) 595-607 (Springer, 2010).

225 3 Vidal, C. *et al.* *Establishing forest inventory reference definitions for forest and growing*  
226 *stock: a study towards common reporting*, Vol. 42 (2008).

227 4 Gschwantner, T. *et al.* Growing stock monitoring by European National Forest  
228 Inventories: Historical origins, current methods and harmonisation. *Forest Ecology*  
229 *and Management* **505**, 119868, doi:<https://doi.org/10.1016/j.foreco.2021.119868>  
230 (2022).

231 5 Köhl, M., Ehrhart, H.-P., Knauf, M. & Neupane, P. R. A viable indicator approach for  
232 assessing sustainable forest management in terms of carbon emissions and removals.  
233 *Ecological Indicators* **111**, 106057, doi:<https://doi.org/10.1016/j.ecolind.2019.106057>  
234 (2020).

235 6 Korosuo, A. *et al.* The role of forests in the EU climate policy: are we on the right track?  
236 *Carbon Balance and Management* **18**, 15, doi:10.1186/s13021-023-00234-0 (2023).


237 7 Vidal, C. *et al.* The role of European National Forest Inventories for international  
238 forestry reporting. *Annals of Forest Science* **73**, 793-806, doi:10.1007/s13595-016-  
239 0545-6 (2016).

- 240 8 Hansen, E., Rahlf, J., Astrup, R. & Gobakken, T. Taper, volume, and bark thickness  
241 models for spruce, pine, and birch in Norway. *Scandinavian Journal of Forest Research*,  
242 1-16, doi:10.1080/02827581.2023.2243821 (2023).
- 243 9 Bouriaud, O., Stefan, G. & Saint-André, L. Comparing local calibration using random  
244 effects estimation and Bayesian calibrations: a case study with a mixed effect stem  
245 profile model. *Annals of Forest Science* **76**, 65, doi:10.1007/s13595-019-0848-5  
246 (2019).
- 247 10 Zell, J., Nitzsche, J., Stadelmann, G. & Thürig, E. SwissStandSim: ein klimasensitives,  
248 einzelbaumbasiertes Waldwachstumsmodell. *Schweizerische Zeitschrift für*  
249 *Forstwesen* **171**, 116-123, doi:10.3188/szf.2020.0116 (2021).
- 250 11 Köhl, M., Magnussen, S. & Marchetti, M. Forest Mensuration in *Sampling Methods,*  
251 *Remote Sensing and GIS Multiresource Forest Inventory* (eds Michael Köhl, Steen  
252 Magnussen, & Marco Marchetti) 17-69 (Springer, Berlin, Heidelberg, 2006).
- 253 12 Zianis, D., Muukkonen, P., Mäkipää, R. & Mencuccini, M. Biomass and stem volume  
254 equations for tree species in Europe. *Silva Fennica Monographs* **4**, 63 (2005).
- 255 13 MacFarlane, D. W. Allometric Scaling of Large Branch Volume in Hardwood Trees in  
256 Michigan, USA: Implications for Aboveground Forest Carbon Stock Inventories. *Forest*  
257 *Science* **57**, 451-459, doi:10.1093/forests/57.6.451 (2011).
- 258 14 Brassel, P. & Lischke, H. *Swiss National Forest Inventory, Methods and models of the*  
259 *second assessment* (Swiss Federal Research Institute WSL, 2001).
- 260 15 Fischer, C. & Traub, B. *Swiss National Forest Inventory – Methods and Models of the*  
261 *Fourth Assessment* (Springer International Publishing, 2019).
- 262 16 Herold, A. *et al.* State and Change of Forest Resources in *Swiss National Forest*  
263 *Inventory – Methods and Models of the Fourth Assessment Managing Forest*  
264 *Ecosystems, vol 35* (eds Christoph Fischer & Berthold Traub) 205-230 (Springer  
265 International Publishing, 2019).
- 266 17 Forrester, D. I., Nitzsche, J. & Schmid, H. The Experimental Forest Management  
267 project: An overview and methodology of the long-term growth and yield plot  
268 network. (Swiss Federal Institute of Forest, Snow and Landscape Research WSL,  
269 Birmensdorf, 2019).
- 270 18 Forrester, D. I., Schmid, H. & Nitzsche, J. The Experimental Forest Management  
271 network. EnviDat, doi:10.16904/envidat.213 (2021).
- 272 19 Kaufmann, E. Estimation of standing timber, growth and cut in *Swiss National Forest*  
273 *Inventory: Methods and Models of the Second Assessment* (eds P. Brassel & H. Lischke)  
274 162–196 (Swiss Federal Research Institute WSL, 2001).
- 275 20 Flury, P. Ergebnisse aus Kahlschlägen in *Mitteilungen der Schweizerischen*  
276 *Centralanstalt für das forstliche Versuchswesen* Vol. 6 (ed C Bourgeois) 87-206 (1898).

- 277 21 Flury, P. Ertragstafeln für die Fichte und Buche der Schweiz. 290 (Zürich, 1907).
- 278 22 Urich, P. Das Urich'sche Holzmassenermittelungs-Verfahren ohne  
279 Kreisflächenberechnung. *Forstwissenschaftliches Centralblatt* **23**, 369-379,  
280 doi:10.1007/BF01841427 (1901).
- 281 23 Fitzpatrick, H. Some methods of estimating the volume of timber in woods and  
282 plantations. *Irish Forestry* **1**, 46-49 (1944).
- 283 24 Gschwantner, T. *et al.* Harmonisation of stem volume estimates in European National  
284 Forest Inventories. *Annals of Forest Science* **76**, 24, doi:10.1007/s13595-019-0800-8  
285 (2019).
- 286 25 Flury, P. Untersuchungen über das spezifische Gewicht des Reisigs in frischgefälltem  
287 Zustande in *Mitteilungen der Schweizerischen Centralanstalt für das forstliche*  
288 *Versuchswesen* Vol. 2 (ed A. Bühler) 15-24 (1892).
- 289 26 Gayer, K. & Fabricius, L. *Die Forstbenutzung ein Lehr- und Handbuch (13. Auflage)*  
290 (Parey, 1935).
- 291 27 Mäkinen, H., Ojansuu, R., Sairanen, P. & Yli-Kojola, H. Predicting branch characteristics  
292 of Norway spruce ( *Picea abies* (L.) Karst.) from simple stand and tree measurements.  
293 *Forestry: An International Journal of Forest Research* **76**, 525-546,  
294 doi:10.1093/forestry/76.5.525 (2003).
- 295 28 Didion, M., Herold, A., Vulovic, Z., Nitzsche, J. & Stillhard, J. Datasets for deriving  
296 functions for the stem- and branchwood volume in the Swiss National Forest  
297 Inventory. doi:<https://www.doi.org/10.16904/envidat.358>. (2022).
- 298 29 Kublin, E., Breidenbach, J. & Kändler, G. A flexible stem taper and volume prediction  
299 method based on mixed-effects B-spline regression. *European Journal of Forest*  
300 *Research* **132**, 983-997, doi:10.1007/s10342-013-0715-0 (2013).
- 301 30 R: A language and environment for statistical computing (R Foundation for Statistical  
302 Computing, Vienna, Austria. URL <https://www.R-project.org/>. Vienna, Austria, 2022).
- 303 31 data.table: Extension of `data.frame`. R package version 1.14.2. [https://CRAN.R-](https://CRAN.R-project.org/package=data.table)  
304 [project.org/package=data.table](https://CRAN.R-project.org/package=data.table) (2021).
- 305 32 Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2016).
- 306 33 `_dplyr: A Grammar of Data Manipulation_`. R package version 1.0.9, <[https://CRAN.R-](https://CRAN.R-project.org/package=dplyr)  
307 [project.org/package=dplyr](https://CRAN.R-project.org/package=dplyr)> (2022).
- 308 34 Didion, M., Herold, A. & Thürig, E. Whole Tree Biomass and Carbon Stock in *Swiss*  
309 *National Forest Inventory – Methods and Models of the Fourth Assessment Managing*  
310 *Forest Ecosystems, vol 35* (eds Christoph Fischer & Berthold Traub) 243-248 (Springer  
311 International Publishing, 2019).


312





314


315 Figure 1. Measurements along the stem. Length below 7 cm diameter (D) from the base of the  
 316 tree, i.e. including stump; section-wise diameter every 2 m along the stem starting at 1m from  
 317 the base of the tree: DM1, DM3, DM5, etc. Additional measurements: Diameter at 0.65 m and  
 318 1.30 m (DBH); mid-diameter and length of the final stem section where  $D \geq 7$  cm if the length  
 319 is  $< 2$  m; and mid-diameter and length of stem top.



321

322 Figure 2. Spatial distribution of the 714 EFM plots with stem and branch data. Note that sites  
 323 may overlap and are not visible and that for 54 plots no detailed spatial information was  
 324 available. The five production regions represent a classification used in the Swiss National  
 325 Forest Inventory indicating relatively homogeneous growth and wood production conditions  
 326 (Glossary in Fischer and Traub<sup>15</sup>). The insert presents the elevation distribution of the plots by  
 327 500 m classes.

328




329

330 Figure 3. DBH distribution in 5cm bins of trees in the current dataset by main tree species (NFI  
 331 classification, cf. Table 1) with corresponding branch volume (coarse and/or fine, i.e. diameter  
 332 threshold of 7 cm) data ('Y') or with stem measurements only ('N'). Note that 24 observations  
 333 with DBH > 1000 mm are not shown.

334

335


a)



336

337

b)



338


339

340

341

Figure 4. Boxplots of the diameter of stem sections at 0.65 m, 1.3 m (i.e., DBH) and starting at 1m every 2 m until the lower threshold of 7 cm is reached, as well as the diameter at half the length of the tree top (i.e, the part of the stem where the it has a diameter of 7 cm and the


342 full height) by main tree species (NFI classification, cf. Table 1) for a) conifers and b)  
343 broadleaves. The values on top of each boxplot give the sample size.



344

345 Figure 5. Volume of coarse branchwood and total of fine woody elements < 7 cm in diameter,  
 346 i.e. including tree top, by main tree species (NFI classification, cf. Table 1). The point  
 347 transparency indicates point density. Sample sizes are given on the right of each panel.

348



349

350 Figure 6. Slenderness ratio (total tree height / DBH) per 10 cm DBH bins by main tree species  
 351 (NFI classification, cf. Table 1).

352

## 353 Tables

Species ID	Species name	NFI main species	N
21	<i>Picea abies</i>	Picea	15'684
22	<i>Abies alba</i>	Abies	7'344
23	<i>Pinus sylvestris</i>	Pinus spp	1'657
24	<i>Larix decidua</i>	Larix	1'629
25	<i>Pinus strobus</i>	Pinus spp	847
26	<i>Pseudotsuga menziesii</i>	other conifers	601
27	<i>Pinus cembra</i>	P. cembra	224
28	<i>Pinus mugo</i> Turra subsp. mugo	Pinus spp	103
29	<i>Picea sitchensis</i>	Picea	29
30	<i>Pinus nigra</i>	Pinus spp	129
31	<i>Abies grandis</i>	Abies	61
32	<i>Chamaecyparis</i>	other conifers	60
33	<i>Cryptomeria japonica</i>	other conifers	21
34	<i>Thuja plicata</i>	other conifers	77
35	<i>Picea omorika</i>	other conifers	14
36	<i>Larix kaempferi</i> (Lamb.) Carrière	Larix	4
41	<i>Fagus sylvatica</i>	Fagus	8'603
42	<i>Quercus petraea</i> , <i>Q. robur</i> , <i>Q. rubra</i>	Quercus	1'821
43	<i>Fraxinus americana</i> , <i>F. excelsior</i>	Fraxinus	153
44	<i>Acer campestre</i> , <i>A. platanoides</i> , <i>A. pseudoplatanus</i>	Acer	96
45	<i>Populus tremula</i>	other broadleaves	216
46	<i>Castanea sativa</i>	Castanea	82
47	<i>Betula pendula</i>	other broadleaves	97
48	<i>Juglans regia</i>	other broadleaves	218
51	<i>Ulmus glabra</i>	other broadleaves	8
52	<i>Prunus avium</i>	other broadleaves	4
60	Other broadleaves, incl. <i>Sorbus</i> spp and <i>Tilia</i> spp	other broadleaves	9

354 Table 1. Tree species information. Species are grouped based on the classification used in the  
355 Swiss National Forest Inventory (Table 14.1 in Didion, et al. <sup>34</sup>). Species was not recorded for  
356 558 trees

357



Variable name	Definition	Value range	N
SiteID	Site descriptor; 8-digit code <sup>17</sup>	01001000 - 62007004	768
Lat*	Latitude of the plot centre [degrees north]	46.08° - 50.57°	714
Long*	Longitude of the plot centre [degrees west]	6.15° - 10.24°	714
Elev*	Elevation [meter above sea level] derived from a digital elevation model	310 - 2000	714
NFI_PR	NFI Production region <sup>15</sup>	Jura, Plateau, Pre- Alps, Alps	768
InvYear	Inventory year	1888 - 1974	40'349
StandAge	Age structure	- even-aged - uneven-aged	33'044 6'727
StandComp	Tree species composition	- pure - conifer mixed (> 75% conifers) - broadleaved mixed (> 75% broadleaves) - conifer-broadleaved mixed	23'685 8'758 620 5'490
TreeID	Running number	1 – 40'349	40'349
TreeSpecies	Species name	See Table 1	39'791
NFI_mainspecies	NFI main species	See Table 1	40'357
TreeAge	age [years]	1 – 43 – 65 – 96 - 340	33'143
DBH	Mean DBH [mm]	6 – 138 – 230 – 341 – 1581	40'349
H_total	Total height [dm]	15 – 151 – 226 -284 - 574	40'349
L_coarsestem	Length of stem from the base to stem D=7 cm [dm] <sup>+</sup>	0– 106 – 192– 252 - 552	40'305
L_coarsestemfinal	Length of the final section of the stem until D=7 cm [dm] if not 2 m in length	0 – 0 – 6 – 10 - 186	40'305
L_top	Length of the tree top (part of the stem where D<7 cm [dm])	2 – 26 – 36 – 46 - 293	40'305
DM065, DM1, DM3, ... DM53	mean stem D at 0.65 m and every 2 m starting at 1m where D >= 7 cm [mm]	Figure 6	Figure 6
D_coarsestemfinal	D of the final section of the stem until D=7 cm [mm] measured at half its length	0 – 0 – 73 – 79 - 175	39'751
D_top	D of the tree top (part of the stem where D<7 cm measured at half its length [mm])	2 – 34 – 39 – 43 – 293	39'571
V_coarsebranch	Volume of coarse branchwood ≥ 7 cm in diameter [dm <sup>3</sup> ]	Figure 5	27'297
V_finewoodytotal	Total volume of fine woody elements < 7 cm in diameter, i.e. including tree top [dm <sup>3</sup> ]	Figure 5	18'980

V_finebranch	Volume fine branchwood < 7 cm in diameter [dm <sup>3</sup> ]	0 – 0 – 5 – 147 - 4210	9'667
--------------	--	------------------------	-------

358 \*54 sites were abandoned after, e.g. clearcutting and have no detailed location information  
359 \*based on trees where the stem diameter at 1 m from the base was  $\geq 7$  cm

360 Table 2. Site (total N=768] and tree specific (total N=40'349) data observed or measured with  
361 units in brackets. For continuous tree data, the value range shows minimum, quartiles, and  
362 maximum. D indicates diameter.

Tree species	Conversion Factor	
	Flury <sup>25</sup>	Badoux
Picea spp.,	0.9	0.9
Abies spp.	0.9	0.9
Pinus spp.	0.9	0.9
Larix	as <i>Picea</i>	0.9
Pseudotsuga menziesii	as <i>Picea</i>	0.9
Other conifers	as <i>Picea</i>	0.9
Fagus spp.	1.0	1.0
Acer spp.	as <i>Fagus</i>	0.9
Alnus spp.	as <i>Fagus</i>	0.9
Betula spp.	as <i>Fagus</i>	0.9
Carpinus spp.	as <i>Fagus</i>	1.0
Fraxinus spp.	0.8	0.8
Populus nigra	as <i>Fagus</i>	0.9
Populus tremula	as <i>Fagus</i>	1.0
Quercus spp.	as <i>Fagus</i>	1.0
Robinia pseudoacacia	as <i>Fagus</i>	0.9
Salix spp.	as <i>Fagus</i>	0.8
Sorbus spp.	as <i>Fagus</i>	0.9
Tilia spp.	as <i>Fagus</i>	0.8
Ulmus	as <i>Fagus</i>	1.0

363 Table 3. Conversion factors based on Flury <sup>25</sup> and E. Badoux (Forest engineer growth and yield,  
364 Federal Institute for Forest Research, predecessor of WSL) to calculate the volume of fine  
365 woody (i.e. diameter < 7 cm) stem and branch material from field measurements of the fresh  
366 weight [kg] of collected standardized bundles of 1m length and 1m circumference [m<sup>3</sup>]. Values  
367 of Badoux were modified from Gayer and Fabricius <sup>26</sup>.

368